

Diabetes Education Centre (DEC)
200 Terrace Hill Street
Brantford, ON, N3R 1G9

Tel: 519-751-5544 ext. 4267 **Fax:** 519-751-5862

Patient Name:	
Date of Birth:	
MD:	
CC:	

Non-Insulin Hypoglycemic Agent Order Set

This set allows DEC Certified Diabetes Educators to make adjustments to current medications using CDA Clinical Practice Guidelines. We will send you a note informing you if we recommend a dose change to a patient. We cannot write prescriptions so please change patient's prescription if you agree to a change made by us.

These orders are to be reviewed annually.

Metformin	□ Reduce by 50% due to significant gastrointestinal side effects.	
(Glucophage)	□ Reduce by 50% if eGFR 30 to 59 mL/minute/1.73m^2.	
	□ Discontinue if eGFR less than 30 mL/minute/1.73m ² .	
Gliclazide MR	Reduce dose by 50% or discontinue due to hypoglycaemia.	
(Diamicron MR)	□ Reduce dose by 50% if eGFR 15 to 29 mL/minute/1.73m^2.	
	□ Discontinue if eGFR less than 15 mL/minute/1.73m^2.	
Glyburide (Diabeta)	Reduce by 50% or discontinue due to hypoglycemia	
	□ Discontinue if eGFR less than 60 mL/minute/1.73m^2.	
GLP-1 agonists	□ Increase as per product monograph to target fasting blood glucose of 5 - 7	
Liraglutide (Victoza)	mmol/L	
Exenatide (Byetta)	Decrease as per product monograph if gastrointestinal side effects.	
	□ Discontinue Liraglutide if eGFR less than 50 mL/minute/1.73m^2	
	□ Reduce Exenatide to to 5mcg bid if eGFR 30 to 59 mL/min/1.73m^2	
	□ Discontinue Exenatide if eGFR less than 30 mL/min/1.73m^2	
eGFR = Estimated Glomerular Filtration Rate CDA=Canadian Diabetes Association		

Physician Name:	
Signature:	Date: